

HYPERSENSPECTRAL OCEAN COLOR SCIENCE: SANTA BARBARA CHANNEL

David A. Siegel
Institute of Computational Earth System Science
University of California at Santa Barbara
Santa Barbara, CA 93106-3060
Phone: 805-893-4547 Fax: 805-893-2578
Email: davey@icess.ucsb.edu
URL: www.icess.ucsb.edu/~davey
Award # N000149710028

LONG-TERM GOALS

Our goal is to develop validation data sets for the modeling and interpretation of hyperspectral satellite ocean color imagery. This grant supports the addition of new optical measurements to the existing Plumes and Blooms (PnB) ocean color observation program. The goal of the PnB *in situ* sampling program is to develop state-of-the-art ocean color algorithms for Case II waters like those found in the Santa Barbara Channel. The PnB project is supported primarily by NASA. This work has filled the most glaring holes in the PnB *in situ* science program. This project provides a fundamental data set for the task of analyzing and modeling hyperspectral reflectance spectra in Case II waters. All PnB data are available via the world wide web (<http://www.icess.ucsb.edu/PnB>).

OBJECTIVES

Our near-term objectives are to validate hyperspectral determinations of remote sensing reflectance, determine *in vivo* absorption spectra for total particulate, dissolved and detrital particulate materials, deploy an *in situ* spectral transmissometer-reflective tube absorption meter, and analyze and interpret these data in conjunction with the rest of the PnB data set. The short-term science goal is to use these data to address the importance of absorption vs. backscattering in the variability of ocean color in Case II waters.

APPROACH

The approach of this grant is to provide supplemental support for the on-going PnB *in situ* science program to provide calibration and validation data sets for the analysis of hyperspectral reflectance spectra. Central to this is the determination of *in vivo* absorption spectra for total particulate ($a_p(\lambda)$), colored dissolved ($a_g(\lambda)$) and detrital particulate materials ($a_{det}(\lambda)$). From these measurements, determinations of the absorption spectrum due to phytoplankton abundances, $a_{ph}(\lambda)$, can be estimated ($a_{ph}(\lambda) = a_p(\lambda) - a_{det}(\lambda)$; figure 1).

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1999		2. REPORT TYPE		3. DATES COVERED 00-00-1999 to 00-00-1999	
4. TITLE AND SUBTITLE Hyperspectral Ocean Color Science: Santa Barbara Channel				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of California at Santa Barbara, Institute of Computational Earth System Science, Santa Barbara, CA, 93106				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified			




Figure 1: *In vivo* phytoplankton (upper left), detrital particulate (upper right), colored dissolved (lower left) and total (lower right) absorption spectra from the PnB program. All observations are shown which were taken between April 1997 and April 1999. The total absorption spectra ($a_{\text{tot}}(\lambda) = a_w(\lambda) + a_{\text{ph}}(\lambda) + a_{\text{det}}(\lambda) + a_g(\lambda)$) where the Pope and Fry [1997] $a_w(\lambda)$ estimates are used.

WORK COMPLETED

To date, we have fully implemented the field aspects of this grant. Since April of 1997, we have collected 357 stations of *in vivo* absorption spectra matched in time with hyperspectral estimates of remote sensing reflectance, $R_{rs}^+(I)$. In addition, as part of every PnB station, samples are collected for inorganic nutrients, HPLC phytoplankton pigment analysis, fluorometric pigment analysis, total suspended solids, and biogenic and lithogenic silica concentrations.

We have also validated hyperspectral estimates of remote sensing reflectance (Toole et al. 1999). We show that the dominant portion of the variability in $R_{rs}^+(I)$ spectra determined using above-water techniques is due to an incomplete removal of sky radiance. We developed a correction procedure for this effect by merging fixed band profiler and above-water hyperspectral estimates of $R_{rs}^+(I)$ spectra (figure 2). The differences are striking and a manuscript on this method has been submitted to *Applied Optics*. As mentioned previously, all PnB data are available via the world wide web (<http://www.icess.ucsb.edu/PnB>).




Figure 2: Before and after sky-radiance correction views of remote sensing reflectance spectra, $R_{rs}^+(I)$, taken from the PnB observational program. All observations are shown which were taken between August 1996 and April 1998. The above-water, hand-held spectrometer technique is used alone in the upper panel and is merged with profiling spectroradiometer data in the lower panel. (Toole *et al.* 1999).

RESULTS

The Plumes and Blooms project has just entered its fourth year, and there are several important results to report. First, we have performed a detailed intercomparison of the three independent methods for determining $R_{rs}^+(I)$ (Toole *et al.* 1998;1999). These results have shown that the above-water hand-held spectrometer technique dramatically overestimates values of $R_{rs}^+(I)$

under moderate to high wind conditions. Comparison with the profiling radiometer and tethered buoy derived $R_{rs}^+(I)$ estimates indicate that the sky correction term required as part of the above-water spectrometer technique is underestimated by a factor of as much as 5. Although the sea surface on the average "sees" the same place in the sky under moderate wind speeds, the collected radiance from the water surface is contaminated not just by the simple reflected beam. This is because the sky radiance distribution is neither geometrically uniform nor slowly varying as a function of viewing angle. That is, the sky correction term cannot be simply modeled using existing procedures (Mueller and Austin, 1995) and improved sky correction procedures are required. A manuscript detailing these analyses and a correction procedure has been submitted for publication to *Applied Optics* (Toole *et al.* 1999).

Second, we have used our observations of $R_{rs}^+(I)$ and $a_{tot}(\lambda)$ from the Santa Barbara Channel to assess the role of spectral backscattering and absorption on ocean color variability in Case II waters. Using the present observations of $R_{rs}^+(I)$ and $a_{tot}(\lambda)$ (figures 1 and 2), estimates of spectral backscatter can be determined (e.g., Garver and Siegel, 1997). We have shown that variability in the reflectance spectra, $R_{rs}^+(I)$, for the Santa Barbara Channel are to first order driven by values of particulate backscatter and to second order by the absorption coefficient, $a_{tot}(\lambda)$. This contrasts blue water sites, such as the Bermuda BioOptics Project (BBOP; Siegel *et al.* 1996), where the changes in the absorption coefficient regulate $R_{rs}^+(I)$ variations in the blue to green portions of the spectrum. We are in the process of writing a manuscript describing the mechanisms of ocean color variations in a Case II environment like the Santa Barbara Channel (Toole *et al.* in prep.).

IMPACTS/APPLICATIONS

First and foremost, the impacts of this project will be in scientific progress. The effects of winds and surface waves on the sky correction term required for the above-water $R_{rs}^+(I)$ estimates limits the utility of this technique to very calm days. Ours is one of the first detailed intercomparison studies of the various $R_{rs}^+(I)$ determinations methods.

Second, the data collected here and the interpretations made using these data mark the scientific basis for developing semi-analytical ocean color algorithms for Case II waters. This work is also on-going and we anticipate significant breakthroughs in ocean color remote sensing capabilities. We have developed objective tuning methodologies for these models and the PnB data set will enable us to assess the importance of region in the design and implementation of ocean color algorithms. Last, all PnB data are available via the world wide web (<http://www.icess.ucsb.edu/PnB>).

TRANSITIONS

The present $R_{rs}^+(I)$ intercomparison results have been presented to the NASA SIMBIOS program and have helped the planning of that program. We have provided code for our semi-analytical ocean color algorithms to Stennis NRL personnel and helped in their implementation

into the SeaDAS system. We have also transitioned some of our algorithms for operational use in SeaWiFS data processing (Siegel et al. 1999).

RELATED PROJECTS

The Bermuda BioOptics Project (BBOP; Siegel *et al.* 1996) is supported by the NASA SIMBIOS program. It provides support for the UCSB calibration laboratory which is used extensively throughout the PnB project.

REFERENCES

- Garver, S.A., and D.A. Siegel, 1997: Inherent optical property inversion of ocean color spectra and its biogeochemical interpretation: I. Time series from the Sargasso Sea. *Journal of Geophysical Research.*, **102**, 18,607-18,625.
- Mueller, J.L., and R.W Austin, 1995: Ocean optics protocols for SeaWiFS Validation, Revision 1. *NASA Tech. Memo. 104566*, Vol. 25, NASA GSFC, Greenbelt, MD, 67 pp.
- Siegel, D.A., M.C. O'Brien, S.A. Garver, E.A. Brody, J.C. Sorensen, A.F. Michaels, E.A. Caporelli and N.B. Nelson, 1996: The Bermuda BioOptics Project - BBOP. *Proc. SPIE, Ocean Optics XIII*, **2963**, 308-313.
- Siegel, D.A., M. Wang, S. Maritorena and W. Robinson, 1999: Atmospheric correction of satellite ocean color imagery: The black pixel assumption. Submitted to *Applied Optics*.
- Toole, D.A., M.J. Neumann, T.K. Westberry, and D.A. Siegel, 1998: Comparison of *in situ* methodologies in Case II environments. In press *Ocean Optics XIV*.
- Toole, D.A., D.A. Siegel, D.W. Menzies, M.J. Neumann and R.C. Smith, 1999: Remote sensing reflectance determinations in coastal environments - Impacts of instrumental characteristics and environmental variability. Submitted to *Applied Optics*.
- Toole, D.A., D.A. Siegel, 1998. Empirical and Mechanistic Approaches to Understanding Ocean Color Variability in the Santa Barbara Channel. ASLO 1999, Santa Fe, New Mexico.
- Toole, D.A., and D.A. Siegel, in prep: Ocean color variability in the Case II waters of the Santa Barbara Channel. For *Continental Shelf Research*.